



**Impatiens SUN
HARMONY (Best in
Quarts and Gallons)**

	pH	Ec	Min Temp F	Max Temp F	Min TEMP C	Max TEMP C	Pinch	Feed	Light	PGR	Fungicide	Comments
Rooting												
Week 1 sticking liner 1 per c	6.0 to 6.5	0.80	72	74	22	23	No Pinch	50 ppm N in mist	Day Neutral		Daconil or Similar chemical day of sticking	
Week 2	6.0 to 6.5	.80 to .90	72	74	22	23	No Pinch	50 ppm N in mist	Day Neutral		Second app Daconil or Similar Chemical	
Week 3	6.0 to 6.5	.80 to .90	72	74	22	23	No Pinch	Feed 65 to 80 ppm N	Day Neutral			
Week 4	6.0 to 6.5	.80 to .90	68	72	20	22	No Pinch	Feed 65 to 80 ppm N	Day Neutral	Bonzi Or Sumagic		
Week 5 Transplant 1 per po	6.0 to 6.5	.80 to .90	68	72	20	22	No Pinch	Feed 65 to 80 ppm N	Day Neutral			
Week 6 to 14	6.0 to 6.5	.80 to .90	68	72	20	22	No Pinch	Feed 65 to 80 ppm N	Day Neutral			Drench Bonzi if needed at 1/4 of 1 ppm

Flores can be sprayed at 300 ppm the day after sticking the unrooted cutting in the liner. This will reset the flower timing. Only 1 hour is required for the mist to be off for the Florel to work.