

Lobelia ANABEL

Habit Recommended Container Flower timing Stick to Flower timing Trailing, Semi aggressive 10 and 12IN Premium Early 12 to 14 weeks

| | рН | Ec | Min Temp F | Max Temp F | Min TEMP C | Max TEMP C | Pinch | Feed | Light | PGR | Fungicide | Comments |
|---------------------------------|------------|------------|------------|------------|------------|------------|-------|-------------|------------|--------------|-----------|------------------------|
| Rooting | · · | | | | | | | | - U | | ŭ | |
| | | | | | | | | | | | Daconil | |
| | | | | | | | | 50 ppm N | | | day of | |
| Week 1 sticking liner | 5.8 to 6.0 | 0.80 | 72 | 74 | 22 | 23 | | in mist | Not Needed | | sticking | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | Second | |
| | | | | | | | | 50 ppm N | | | арр | |
| Week 2 | 5.8 to 6.0 | .80 to .90 | 72 | 74 | 22 | 23 | | in mist | Not Needed | | Daconil | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| l | | | | | | | | Feed 100 | | B-9/Arest | | B-9 3750 ppm / Arest |
| Week 3 | 5.8 to 6.0 | .90 to 1.2 | 65 | 72 | 18 | 22 | Pinch | to 150 ppm | Not Needed | if needed | | 5ppm tank mix |
| | | | | | | | | | | | | |
| | | | | | | | | feed 100 to | | B- 9 / Arest | | |
| Week 4 | 5.8 to 6.0 | .90 to 1.2 | 65 | 72 | 18 | 22 | | | Not Needed | | | B-9 3750 ppm if needed |
| vveek 4 | 5.0 10 0.0 | .90 to 1.2 | 65 | 12 | 10 | 22 | | 150 ppiii | NOT Needed | ii needed | | b-9 3750 ppm ii needed |
| | | 1 | | | | | | feed 100 to | | | | |
| Week 5 Transplant 3 per basket | 5.8 to 6.0 | .90 to 1.2 | 65 | 72 | 18 | 22 | | | Not Needed | | | |
| Trock o Transplant o per basket | 3.0 10 0.0 | .55 10 1.2 | 33 | | | | | .cc ppiii | | | | |
| | | | | | | | | feed 100 to | | | | Bonzi Drench 3ppm |
| Week 6 to 12 or 14 | 5.8 to 6.0 | .90 to 1.2 | 65 | 72 | 18 | 22 | | | Not Needed | | | when to side of pot |

Bonzi is effective as a drench Read label before use