

# HARMONY<sup>®</sup>

New Guinea Impatiens



**HARMONY<sup>®</sup> Bicolor Red**



- Top selling NGI series in North America
- Easy to grow and very reliable in production
- Great in 12 cm/quarts up to hanging baskets
- Mature height and width is 16" x 12" (41cm x 30cm)

Our best-selling varieties in the HARMONY<sup>®</sup> series:



**HARMONY<sup>®</sup>  
Violet**



**HARMONY<sup>®</sup>  
Orange Blaze**



**HARMONY<sup>®</sup>  
Snow**



**HARMONY<sup>®</sup>  
Magenta**



**HARMONY<sup>®</sup>  
Red Cardinal**



**HARMONY<sup>®</sup>  
Dark Lilac**



**HARMONY<sup>®</sup>  
Candy Cream**



**HARMONY<sup>®</sup>  
Purple Cream**



**HARMONY<sup>®</sup>  
Tangerine**



**HARMONY<sup>®</sup>  
Fire**

For additional technical growing information:

[HARMONY<sup>™</sup> CLICK HERE](#)



 **DANZIGER**

[www.danzigeronline.com](http://www.danzigeronline.com) |   

# HARMONY<sup>®</sup>

New Guinea Impatiens



## PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer  
Flowering



Width

(12 - 14")  
30 - 36 cm



Height

(14 - 18")  
36 - 46 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21°-22° C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68°-70° F (20°-21° C)		Weeks 3-5 80 ppm N	Apply broad spectrum Week 2



- Spray adjuvant at sticking



- Spray Ethephon at 300 ppm the day after stick. This will reset flowering response. One hour of Ethephon on the leaves will be sufficient time while mist if off.
- Maintain warm temperatures allowing plants to develop and flower
- Average daily temperature of 70 F

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	6.0 / 6.5	0.50	72°-74° F (22°-23° C)		Not required		Day of sticking	
Week 2	6.0 / 6.5	.80 to .90	72°-74° F (22°-23° C)	50 ppm N in mist	Not required		Second fungicide app	
Week 3	6.0 / 6.5	.80 to .90	72°-74° F (22°-23° C)	80 ppm N in mist	Not required			
Week 4	6.0 / 6.5	.80 to .90	70°-72° F (21°-22° C)	80 ppm N in mist	Not required			
Week 5	6.0 / 6.5	.80 to .90	70°-72° F (21°-22° C)	80 ppm N in mist	Not required			Pinch is not recommended

For additional technical growing information:

HARMONY™ [CLICK HERE](#)



 **DANZIGER**  
www.danzigeronline.com |   

# HARMONY<sup>®</sup>

## New Guinea Impatiens



### FINISHING TIPS

- Maintain a low EC level for proper growth and sizing of plant
- Transplant liners when plants are rooted and actively growing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
7 to 8 wks (12 cm) 5" pot	Average Day 68°-70° F (20°-21° C)	Pinch –No  Daylength Neutral	100 - 125 ppm N	If required, Uniconazole at 5 to 7 ppm spray application early in crop- Paclobutrazol at 0.5 ppm drench within 3 weeks of finishing
8 to 9 wks (15 cm) 6 ½"	68°-70° F (20°-21° C)		Soil EC 0.8 - 1.0 pH 6.0 to 6.5	

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Uniconazole 5 to 7 ppm early in crop  
Paclobutrazol drench application at 0.5 ppm for finishing
- Spray broad spectrum fungicide after transplant
- Scout for Aphids and Thrips on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.8 to 6.2	1.0 to 1.2	65°-68° F (18°-20° C)	feed 100 to 150 ppm		Uniconazole spray	Drench fungicide at transplant	
<b>Week 6 to 13</b>	5.8 to 6.2	1.0 to 1.2	65°-68° F (18°-20° C)	feed 100 to 150 ppm		Paclobutrazol Drench at 0.5 ppm		Allow plants to reach 85% of required size before PGR application

For additional technical growing information:

HARMONY™ [CLICK HERE](#)

