

PAZZAZ NANO™

Portulaca oleracea



BRAND NEW 23



PAZZAZ™ NANO
Orange Twist

- Naturally compact, trailing habit that works well for small and mid-sized containers
- Perfect for heat tolerant/drought tolerant programs
- Available in a wide array of vibrant colors, including bicolors
- Mature height and width is 8 inches x 12 inches (20cm x 30cm)

More varieties in the PAZZAZ NANO™ series:



PAZZAZ™ NANO
Yellow Twist



PAZZAZ™ NANO
Gold



PAZZAZ™ NANO
Tropical Punch



PAZZAZ™ NANO
Fuchsia Improved



PAZZAZ NANO™
Purple



PAZZAZ NANO™
Candy Pink



PAZZAZ NANO™
Fuchsia



PAZZAZ NANO™
Yellow



PAZZAZ NANO™
Hot Pink



PAZZAZ NANO™
Orange

For additional technical growing information:

PAZZAZ NANO™ [CLICK HERE](#)



 **DANZIGER**
www.danzigeronline.com |   

PAZZAZ NANO™

Portulaca oleracea



PROPAGATION TIPS

- Stick on priority – Number 1 out of 4 categories



Flowering time

Late Spring / Summer
Flowering



Width

(10 - 12")
25- 30 cm



Height

(8 - 10")
20 - 25 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21°-22° C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68°-70° F (20°-21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking. Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72°-74° F (22°-23° C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72°-74° F (22°-23° C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68°-70° F (20°-21° C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	68°-70° F (20°-21° C)	feed 100 to 150 ppm		Daminozide as required		

For additional technical growing information:

PAZZAZ NANO™ [CLICK HERE](#)



 **DANZIGER**
www.danzigeronline.com |   

PAZZAZ NANO™

Portulaca oleracea



FINISHING TIPS

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks (12 cm) 5" pot	Average Day 68°-72° F (20°-22° C)	Pinch Additional lighting will hasten flowering under short days	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing if required
6 to 7 wks (15 cm) 6 ½"			Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
7 to 9 wks (20 cm) 8"	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide control height
- Scout for Aphids, Thrips and Whiteflies

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.8 to 6.2	1.0 to 1.2	68°-72° F (20°-22° C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	
Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68°-72° F (20°-22° C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm for finishing		

For additional technical growing information:

PAZZAZ NANO™ [CLICK HERE](#)

