


LUCILLE™

Leucanthemum superbum



LUCILLE™ Grace

Best for 2G / 30-36 cm pots and larger

- Cold Hardiness 4 - 9
- No vernalization or cooling required (First Year Flowering)
- Early spring flower response
- Lucille Chic has a compact habit, Lucille White has a medium habit and Lucille Grace has the most vigor and largest flowers
- Mature height and width is 46 cm x 36 cm (18 x 14 inches)

More varieties in the LUCILLE™ series:



LUCILLE™ White

Best for Gallons
/ 15-18 cm pots



LUCILLE™ Chic

Best for Quarts and
Gallons / 12-18 cm
pots

For additional technical growing information:

LUCILLE™ [CLICK HERE](#)



LUCILLE™

Leucanthemum superbum



PROPAGATION TIPS

- Stick on priority – Number 3 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

36 - 41 cm
(14 - 16")



Height

30 - 51 cm
(12 - 20")

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Suggested	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 18° - 20° C (65° - 68° F)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2



Irrigation specification – Spray with adjuvant the day of sticking/Remove from mist as soon as possible.



K IBA spray application will hasten and even rooting. Allow soil to dry slightly between irrigation

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.0	0.80	21 / 22 C	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	21 / 22 C	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.80 to .90	Cool to 20 / 21 C	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	.80 to 1.2	20 to 21 C	Feed 100 to 150 ppm		Daminozide as needed		
Week 5	5.8 to 6.2	.80 to 1.2	18 to 20 C	Feed 100 to 150 ppm		Daminozide as needed		Pinch is not required

For additional technical growing information:

LUCILLE™ [CLICK HERE](#)



DANZIGER

www.danzigeronline.com |

LUCILLE™

Leucanthemum superbum



FINISHING TIPS

- Maintain soil moisture evenly
- High light levels are best for finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
9 to 10 wks	Average Day	Pinch –None required	100 - 150 ppm N	Daminozide spray application @ 2500 – 3750 ppm
15 cm (1 gallon)	(18°C)			
10 to 11 wks			Soil EC 1.0 - 1.5 pH 5.8 to 6.2	
20 cm (2 gallon)				

		Recommended Chemicals
Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide at 2000 to 3750 ppm on finished plants early in crop schedule
- Maintain good airflow and allow plants to dry before nightfall
- Scout for Aphids, Leafminors, Mites and Thrips
- Drench after transplant for Botrytis and Pythium / Phytophthora

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	1.0 to 1.2	18° to 20°C	feed 150 to 200 ppm			Drench fungicide after transplant	Control of Pythium & Phytophthora
Week 6 to 14	5.8 to 6.2	1.0 to 1.5	18° to 20°C	feed 150 to 200 ppm		Daminozide as needed		

For additional technical growing information:

LUCILLE™ [CLICK HERE](#)

