

'PINK PEARL' PPAF

TERRA NOVA
NURSERIES, INC.

'PARADE OF PARROTS' PPAF

Agastache hybrida

- First year flowering, no vernalization or bulking required
- Early and uniform bloom
- Re-blooms for color all summer long
- Strong radial branching



Upright



45-50 cm
18-20 in



50-55 cm
20-22 in



Zone
6-10



15-18
Gal+



Landscape



Pink pearl

'PINK PEARL' PPAF

'PARADE OF PARROTS' PPAF



Pink Pearl *PPAF*



Parade of parrots *PPAF*

PINK PEARL & PARADE OF PARROTS

Agastache Hybrida

For additional information and availability:

[PINK PEARL](#)

[PARADE OF PARROTS](#)



Stick on priority – Number 2 out of 4 categories

PROPAGATION TIPS

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-4 100 to 125 ppm	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6/6.2	0.80	70°/72°F (21°/22°C)	50 ppm N in mist	Optional		Day of sticking	
Week 2	5.6 to 6.2	.80 to .90	70°/72°F (21°/22°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.2	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm N				
Week 4	5.6 to 6.2	.90 to 1.0	68° to 70°F (20° to 21°C)	Feed 100 ppm N				
Week 5	5.6 to 6.2	.90 to 1.0	68° to 70°F (20° to 21°C)	Feed 100 ppm N		Optional		



– **Irrigation specification** – Spray with adjuvant the day of sticking/Remove from mist as soon as possible. Avoid saturated soil conditions.



Daminozide spray application @ 1500 ppm week 4 may be required to control stretch in liner stage. Once rooted, allow soil to dry slightly between irrigations

PINK PEARL & PARADE OF PARROTS

Agastache Hybrida

For additional information and availability:

[PINK PEARL](#)

[PARADE OF PARROTS](#)

- High quality plants are finished with high light and cool temperatures
- Pinch at transplant
- Allow some growth under short days to 'bulk plants before initiation

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 10 wks	Average Day	Pinch required/ long day flowering response	100 - 150 ppm N	Daminozide spray at 2,000 to 2,500 ppm
15 cm (1 gallon)	(16°-18° C)			
10 to 12 wks	1-2 plants per container		Soil EC 0.8 - 1.0 pH 5.6 to 6.5	
20 cm (2 gallon)				

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium	PROPAMOCARB / MEFENOXAM
	Phytophthora	
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB / Triflumizole

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.2	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm			Drench fungicide after transplant	Control of Pythium & Phytophthora
Week 6 to 12	5.6 to 6.2	0.8 to 1.0	60° to 65° F (16° to 18° C)	feed 150 to 200 ppm		Daminozide spray at 2,000 to 2,500 ppm	Spray fungicide in finishing protocols	Control of Bacterial Leaf Blights



- Facultative Long Day Flowering Response – June through September
- Natural foliar disease resistance
- Maintain good airflow and allow plants to dry between irrigations
- Drench after transplant for Rhizoctonia and Pythium / Phytophthora