


SPLASH DANCE™

Petunia hybrida



- Selected to hold its speckled pattern into the heat
- Mounded habit works well in baskets and combos
- Blooms early enough for any market
- Mature Height and Width 18 inches X 24 inches (46cm x 61cm)

SPLASH DANCE™
Moon Walk

More varieties in the SPLASH DANCE™ series:



SPLASH DANCE™
Fuchsia Flamenco



SPLASH DANCE™
Rumba Rose



SPLASH DANCE™
Violet Vogue



SPLASH DANCE™
Poppin' Pink



SPLASH DANCE™
Bolero Blue



SPLASH DANCE™
Magenta Mambo



SPLASH DANCE™
Purple Polka

For additional technical growing information:

[SPLASH DANCE™ CLICK HERE](#)



 **DANZIGER**
www.danzigeronline.com |   

SPLASH DANCE™

Petunia hybrida



PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

(18 - 24")
46 - 61 cm



Height

(18 - 24")
46 - 61 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification – Reduce mist as soon as possible



The series is well branched so pinching is not need for a quality finish. Daminozide at 1500 ppm the day after sticking. Week 4 – Daminozide @ 2500 ppm spray application

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.0	0.80	72 / 74 F	50 ppm N in mist	Additional lighting is optional depending on region	Daminozide day after sticking	Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	72 / 74 F	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.0	.90 to 1.0	Cool to 68 F	Feed 100 ppm to 150 ppm				
Week 4	5.6 to 6.0	1.0 to 1.2	65 to 68 F	feed 100 to 150 ppm		Daminozide@ 2500 spray application		

For additional technical growing information:

SPLASH DANCE™ [CLICK HERE](#)



SPLASH DANCE™

Petunia hybrida



FINISHING TIPS

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 68° to 72°F (20° to 22°C)	No pinch necessary	150 - 200 ppm N	Finish stages Daminozide @ 2500 to 3200 ppm
8 to 9 wks (15 cm) 6 ½"		Additional lighting is optional depending on region	Soil EC 1.2 - 1.5 pH 5.6 to 6.0	
8 to 10 wks (20 cm) 8"	3 plants per container			
8 to 10 wks (25 cm) 10"	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- PGR - Spray Daminozide at 2,500 to 3,200 ppm and / or Paclobutrazol drench application at 1 to 2 ppm for finishing if required. The most vigorous varieties are Bolero Blue, Purple Polka and Violet Vogue

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.6 to 6.0	1.0 to 1.2	65 to 68 f	feed 100 to 150 ppm	N/A	Daminozide/ Uniconazole tank mix if required	Spray fungicide after transplant	
Week 5 to 15	5.6 to 6.0	1.2 to 1.5	65 to 68 f	feed 100 to 150 ppm	N/A	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

For additional technical growing information:

SPLASH DANCE™ [CLICK HERE](#)

