

# PAZZAZ™

Portulaca oleracea



BRAND  
NEW 23



**PAZZAZ™ Red Flare Improved**

- Moderate trailing habit that works well for gallon and hanging basket production
- Perfect for heat tolerant/drought tolerant programs
- Available in a wide array of vibrant colors
- Mature height and width is 12 inches x 14 inches (30cm x 36cm)

More varieties in the PAZZAZ™ series:



**PAZZAZ™ Fuchsia**



**PAZZAZ™ Tangerine**



**PAZZAZ™ White**



**PAZZAZ™ Yellow imp.**

For additional technical growing information:

[PAZZAZ™ CLICK HERE](#)



# PAZZAZ™

Portulaca oleracea



## PROPAGATION TIPS

- Stick on priority – Number 1 out of 4 categories



Flowering time

Late Spring / Summer  
Flowering



Width

(14 - 16")  
36- 41 cm



Height

(10 - 12")  
25 - 30 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



**Irrigation specification** – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking. Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68 / 70 F	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	68 to 70 F	feed 100 to 150 ppm		Daminozide as required		

For additional technical growing information:

PAZZAZ™ [CLICK HERE](#)



# PAZZAZ™

Portulaca oleracea



## FINISHING TIPS

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks (12 cm) 5" pot	Average Day 68° to 72°F (20° to 22°C)	Pinch  Additional lighting will hasten flowering under short days	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing
6 to 7 wks (15 cm) 6 ½"			Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
7 to 9 wks				
(20 cm) 8"	3 plants per container			

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide control height
- Scout for Aphids, Thrips and Whiteflies

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 4 Transplant</b>	5.8 to 6.2	1.0 to 1.2	68 to 72 f	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	
<b>Week 5 to 12</b>	5.8 to 6.2	1.2 to 1.5	68 to 72 f	feed 100 to 150 ppm		Daminozide @ 2500 ppm for finishing		

For additional technical growing information:

PAZZAZ™ [CLICK HERE](#)

