


SCOPIA® GULLIVER™

Bacopa / Sutera



- Large flower size
- Well branched genetics
- Full spectrum of colors
- Great product for combinations

SCOPIA® GULLIVER™
Pink

More varieties in the GULLIVER™ series:



SCOPIA® GULLIVER™
White Improved



Best Seller

SCOPIA® GULLIVER™
Dynamic White



Best Seller

SCOPIA® GULLIVER™
Blue



SCOPIA® GULLIVER™
Pink Heart Improved

For additional technical growing information:

SCOPIA® GULLIVER™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   

SCOPIA® GULLIVER™

Bacopa / Sutera



PROPAGATION TIPS

- Stick on priority – Number 1 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

(15 - 17")
35 - 40 cm



Height

(4 - 6")
10 - 15 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification - Remove from mist as soon as possible to avoid leaf disease issues.



Daminozide at 1500 ppm day of sticking. The Scopia Gulliver™ Series is well branched so no pinch should be required. Spray fungicide to control leaf diseases.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist		Day of sticking	Day of sticking	
Week 2	5.6 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.2	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm to 150 ppm				
Week 4	5.6 to 6.2	1.0 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm		Daminozide if needed		

For additional technical growing information:

SCOPIA® GULLIVER™ [CLICK HERE](#)



SCOPIA® GULLIVER™

Bacopa / Sutera



FINISHING TIPS

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Consistent soil moisture is best to maximize root structure and plant health
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 65° - 68° (18° - 20° C)	Pinch / Optional	100 - 150 ppm N	Spray application of Daminozide if required
8 to 9 wks (15 cm) 6 ½"	65° - 68° (18° - 20° C)	No lighting necessary	Soil EC 0.8 - 1.0 pH 5.6 to 6.2	
8 to 9 wks (20 cm) 8"	3 plants per container			
8 to 9 wks (25 cm) 10"	3 plants per container			

Pests	Diseases
Aphids	Botrytis
Thrips	Pythium
	Phytophthora



- Consistent soil moisture levels are best – allow soil to dry slightly between irrigations
- Fungicide applications and good air movement to reduce disease pressures
- Scout for Aphids on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.0	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Day Neutral	Daminozide spray application if needed	Spray fungicide after transplant	
Week 6 to 12	5.6 to 6.0	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Not Required			

For additional technical growing information:

SCOPIA® GULLIVER™ [CLICK HERE](#)

