


PAVILION™

Caryopteris incana



- Cold Hardiness 6 – 10
- Short day plant – flowering begins week 36
- No vernalization or cooling required (First Year Flowering)
- Mature height and width is 51 cm x 41 cm (20 x 16 inches)

PAVILION™
Pink

More varieties in the PAVILION™ series:



PAVILION™
White



PAVILION™
Blue

For additional technical growing information:

PAVILION™ [CLICK HERE](#)



PAVILION™

Caryopteris incana



PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time
Spring / Summer
Flowering



Width
41 cm
(16")



Height
51 cm
(20")

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Suggested	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 18° - 20° C (65° - 68° F)	PAVILION	Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2

 **Irrigation specification** – Spray with adjuvant the day of sticking/Remove from mist as soon as possible

 **Caryopteris incana** respond to short days for flowering. Stick roughly week 26 for finishing in the fall. Daminozide at 1500 to 2500 ppm week 4 may be required to control stretch in liner stage.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.5 / 6.2	0.80	70 / 72 F (21 / 22 C)	50 ppm N in mist			Day of sticking	
Week 2	5.5 / 6.2	.80 to .90	70 / 72 F (21 / 22 C)	50 ppm N in mist	Optional		Second fungicide app	
Week 3	5.5 / 6.2	.90 to 1.0	Cool to 65 / 68 F (18 / 20 C)	Feed 100 ppm to 150 ppm	Optional	Daminozide as needed		
Week 4	5.5 / 6.2	1.0 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Optional	Daminozide as needed		Pinch at transplant

For additional technical growing information:

PAVILION™ [CLICK HERE](#)



PAVILION™

Caryopteris incana



FINISHING TIPS

- Pinching is required for branching and controlling height
- High quality plants will finish in all regions / hot and cool regions

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 10 wks	Average Day	Pinch – Suggested day of transplant	100 - 150 ppm N	Daminozide spray application @ 2000 – 2500 ppm
15 cm (1 gallon)	(20° to 21°C)			
8 to 10 wks			Soil EC 1.0 - 1.2 pH 5.5 to 6.2	
20 cm (2 gallon)				

		Recommended Chemicals
Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB
	Rhizoctonia	Pentachloronitrobenzene, Triflumizole



- PGR - Spray Daminozide at 2000 to 2500 ppm on finished plants early in crop schedule
- Maintain good airflow and allow plants to dry before nightfall. Do not keep consistently wet or root rot problems may develop.
- Scout for Aphids, Mites and Whiteflies
- Drench after transplant for Pythium / Phytophthora and Rhizoctonia

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21°C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	Control of Pythium & Rhizoctonia
Week 6 to 15	5.6 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21°C)	feed 100 to 150 ppm				

For additional technical growing information:

PAVILION™ [CLICK HERE](#)

