


# GUATEMALA™

Echinacea hybrid



**Guatemala™ Papaya**



- Cold Hardiness 5 - 9
- No vernalization or cooling required (First Year Flowering)
- Natural June flowering response
- Second vegetative Echinacea available from URC
- Mature height and width is 20 cm x 36 cm (8 x 14 inches)



**Guatemala™ Gold**

For additional technical growing information:

GUATEMALA™ [CLICK HERE](#)



# GUATEMALA™

Echinacea hybrid



## PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer  
Flowering



Width

18 - 26 cm  
(7 - 10")



Height

30 - 38 cm  
(12 - 15")

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 18° - 20° C (65° - 68° F)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2



**Irrigation specification** – Spray with adjuvant the day of sticking/Remove from mist as soon as possible



K-IBA spray application will hasten and even rooting response. Echinacea Panama™ are long day flowering with the natural flowering response being June (facultative long day plant). An adequate substrate for young plants with a pH of 5.8 to 6,5 and a EC of 0,8 mS.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.0	0.80	70 / 72 F (21 / 22 C)	50 ppm N in mist	Required 12 hr daylength		Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	70 / 72 F (21 / 22 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	.90 to 1.0	68 to 70 F (20 to 21 C)	Feed 100 to 150 ppm				
Week 5	5.8 to 6.2	.90 to 1.0	68 to 70 F (20 to 21 C)	Feed 100 to 150 ppm		Daminozide if needed		1500 to 2000 ppm Daminozide

For additional technical growing information:

GUATEMALA™ [CLICK HERE](#)



# GUATEMALA™

Echinacea hybrid



## FINISHING TIPS

- No pinch should be needed
- Naturally compact plant habit
- Spray for bacterial leaf blights

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 10 wks	Average Day	Pinch not required/ extend daylength to 12 hours for early flowering	100 - 150 ppm N	Daminozide spray application @ 2000 – 2500 ppm
15 cm (1 gallon)	(20° to 21°C)			
8 to 10 wks	3 plants per container		Soil EC 0.8 – 1.2	
20 cm (2 gallon)			pH 5.8 to 6.2	

		Recommended Chemicals
Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Long Day Flowering Response Natural flowering time is weeks 24 - 27
- PGR - Spray Daminozide at 2000 to 2500 ppm on finished plants early in crop schedule if needed
- Maintain good airflow and allow plants to dry before nightfall/spray for bacterial leaf blights
- Drench after transplant for Rhizoctonia and Pythium / Phytophthora

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 6 Transplant</b>	5.8 to 6.2	0.8 to 1.0	68° to 70° F (20° to 21°C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	Control of Pythium & Phytophthora
<b>Week 6 to 13</b>	5.8 to 6.2	0.8 to 1.2	68° to 70° F (20° to 21°C)	feed 150 to 200 ppm			Spray fungicide in finishing protocols	Control of Bacterial Leaf Blights

For additional technical growing information:

GUATEMALA™ [CLICK HERE](#)



# GUATEMALA™

Echinacea hybrid



## Israel November 2022



For additional technical growing information:

GUATEMALA™ [CLICK HERE](#)

