

# MUSICA®

Impatiens walleriana



Best Seller

**MUSICA©  
Pink Aroma**



- Spectacular double flowers in a wide range of colors
- Uniform compact growth habit across the series
- Naturally branched so no pinch necessary
- Great in 12 cm/quarts up to hanging baskets
- Mature height and width is 13" x 18" (33 cm x 46 cm)

More varieties in the MUSICA© series:



**MUSICA©  
Lavender**



**MUSICA©  
Bicolor Red**



**MUSICA©  
Elegant Red**



**MUSICA©  
Bicolor Pink**



**MUSICA©  
Electric Purple**

For additional technical growing information:

[MUSICA© CLICK HERE](#)



# MUSICA®

Impatiens walleriana



## PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer  
Flowering



Width

(15 - 20")  
35 - 45 cm



Height

(10 - 15")  
25 - 35 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-5 80 ppm N	Apply broad spectrum Week 2



- Spray adjuvant at sticking. Remove from mist as soon as possible



- Shade will be required for later spring/summer production
- Maintain warm temperatures allowing plants to develop
- Average daily temperature of 70 F
- Spray fungicide to control botrytis

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist			Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.80 to .90	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm N				
Week 4	5.8 to 6.2	.80 to .90	68 to 70 F (20 to 21 C)	Feed 100 ppm N		Should not be required		

For additional technical growing information:

MUSICA© [CLICK HERE](#)



# MUSICA®

Impatiens walleriana



## FINISHING TIPS

- Shade will be required for later spring/summer production
- Maintain a low EC level for proper growth and sizing of plant
- Transplant liners when plants are rooted and actively growing

Average Time (from liners)	Temperature	Pinch/Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 68° - 70° (20° - 21° C)	Pinch –No Daylength Neutral	100 - 125 ppm N	Paclobutrazol at 0.5 ppm drench within 3 weeks of finishing. Allowing soil to dry between irrigations and modest fertility will keep plants more compact
8 to 9 wks (15 cm) 6 ½"	68° - 70° (20° - 21° C)		Soil EC 0.8 - 1.0 pH 5.8 to 6.2	

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Spray broad spectrum fungicide after transplant
- Scout and spray for Powdery Mildew
- Scout for Aphids Leafminers and Thrips on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.8 to 6.2	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 125 ppm N	Day neutral	Paclobutrazol at 0.5 ppm drench within 3 weeks of finishing	Drench fungicide at transplant	
<b>Week 6 to 13</b>	5.8 to 6.2	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 125 ppm N				

For additional technical growing information:

MUSICA© [CLICK HERE](#)

