


STREAM™ Compact

Lobularia



**STREAM™
Compact White**



- Good heat tolerance and resistance to cold
- Flowers from early spring until frost
- Makes an excellent landscape plant
- Great for attracting pollinators

More varieties in the STREAM™ Compact series:



**STREAM™
Compact Rose**



**STREAM™
Compact Purple**




**STREAM™
Compact Violet**

For additional technical growing information:

STREAM™ Compact [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   

STREAM™ Compact

Lobularia



PROPAGATION TIPS

- Stick on priority – Number 1 out of 4 categories



Flowering time
Spring / Summer
Flowering



Width
(4 -6")
10 - 15 cm



Height
(10 - 18")
24 - 46 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification - Remove from mist as soon as possible



Application of K-IBA can hasten rooting. Daminozide at 1500 to 2500 ppm day of sticking. Lighting in propagation is not necessary. The Lobularia Stream™ Series is well branched and a pinch would only be necessary if grown under lower light levels.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.2 / 5.8	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist			Day of sticking	Feeding through Mist
Week 2	5.2 to 5.8	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.2 to 5.8	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.2 to 5.8	1.0 to 1.2	65 to 68 F (18 / 20 C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 5	5.2 to 5.8	1.0 to 1.2	65 to 68 F (18 / 20 C)	Feed 100 ppm to 150 ppm		Daminozide as needed		

For additional technical growing information:

STREAM™ Compact [CLICK HERE](#)



www.danzigeronline.com |

STREAM™ Compact

Lobularia



FINISHING TIPS

- Pinch is optional
- Cool temperature finishing for best quality

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 65° - 68° (18° - 20° C)	Pinch / Optional	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½"	65° - 68° (18° - 20° C)		Soil EC 0.8 - 1.0 pH 5.6 to 6.0	
8 to 9 wks (20 cm) 8"				
8 to 9 wks (25 cm) 10"	3 plants per container			
		No lighting necessary		
	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Consistent soil moisture levels are best – allow soil to dry slightly between irrigations
- Scout for Aphids on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.0	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
Week 6 to 12	5.6 to 6.0	0.8 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

For additional technical growing information:

STREAM™ Compact [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   