

PRIMAVERA™

Pericallis hybrida



**Primavera™
Fuchsia**

- Early flowering time
- Suitable for small pots
- Compact with mounded habit
- Good heat tolerance in the Pericallis category
- Finished height and width is 12 inches x 12 inches (30cm x 30cm)

Other top selling varieties in the PRIMAVERA™ series:



**Primavera™
Denim**



**Primavera™
Blueberry**



**Primavera™
Lavender**



**Primavera™
Navy Blue**

For additional technical growing information:

PRIMAVERA™ [CLICK HERE](#)



PRIMAVERA™

Pericallis hybrida



PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

(12 - 14")
30 - 36 cm



Height

(12 - 14")
30 - 36 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Recommended	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 ppm N	Spray fungicide to control Botrytis and bacteria Week 2



Irrigation specification – Spray adjuvant after sticking. Do not over-mist or saturate soil.



Drench plants with fungicide to protect against root and stem pathogens. Daminozide at 2500 ppm if required week 4 / 5. Pinch the week before transplant or at transplant. Provide adequate air-movement and allow plants to dry going into the evening.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist		Daminozide day of sticking	Day of sticking	Feeding through Mist
Week 2	5.6 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.2	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm				
Week 4	5.6 to 6.2	1.0 to 1.2	65 to 68 F (18 to 20 C)	Feed 100 ppm		Daminozide as needed		
Week 5	5.6 to 6.2	1.0 to 1.2	65 to 68 F (18 to 20 C)	Feed 100 ppm		Daminozide as needed		

For additional technical growing information:

PRIMAVERA™ [CLICK HERE](#)



PRIMAVERA™

Pericallis hybrida



FINISHING TIPS

- Spray fungicide to protect against Powdery Mildew – Supply adequate airflow
- Cool temperature finishing for best quality – 55 to 60 F average daily temperature

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
7 to 8 wks (12 cm) 5" pot	Average Day 55° - 60° (13° - 16° C)	Pinch week before transplant or at transplant	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½"	1 plant per container	Day Neutral	Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
8 to 9 wks (20 cm) 8"		3 plants per container		

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIACARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Allow soil to dry between irrigations / For southern regions, some shade may be necessary
- Best to feed with Nitrate forms of Nitrogen
- Daminozide at 2500 to 3750 ppm / Paclobutrazol at 1 to 2 ppm to finish crop
- Fuchsia, Navy Blue and Violet are the earliest to flower and most vigorous
- Scout for Aphids, Mites and Whiteflies

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	1.0 to 1.2	55 to 60 F (13 to 15 C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	
Week 6 to 12	5.8 to 6.2	1.0 to 1.2	55 to 60 F (13 to 15 C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm	Control of water molds and Thielaviopsis	Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

For additional technical growing information:

PRIMAVERA™ [CLICK HERE](#)

