


SALLYFUN™

Salvia farinacea



Blue Lagoon



- Fast growth rate and early bloomer
- Full plant with upright habit
- Strong flower pedicle
- Flower timing matched across the colors

More varieties in the SALLYFUN™ series:

Medium



SALLYFUN™
Bicolor Blue



SALLYFUN™
Blue Ice



SALLYFUN™
Blue



SALLYFUN™
Deep Ocean



SALLYFUN™
Pure White

Large



SALLYFUN™
Blue Emotion



SALLYFUN™
Sky Blue



SALLYFUN™
Snowwhite

For additional technical growing information:

SALLYFUN™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   




SALLYFUN™

Salvia farinacea



PROPAGATION TIPS

- Stick on priority – Number 3 out of 4 categories

		
Flowering time	Width	Height
Spring / Summer Flowering	(10 - 14") 25- 36 cm	(14 - 20") 36 - 46 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65° - 70° (18° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification – Reduce mist as soon as possible



Spray adjuvant at sticking . Daminozide at 2000 ppm the day after sticking.
Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist	Neutral	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	65 to 68 F (18 / 20 C)	feed 100 to 150 ppm		Daminozide as required		

For additional technical growing information:

SALLYFUN™ [CLICK HERE](#)



SALLYFUN™

Salvia farinacea



FINISHING TIPS

- Allow plants to dry between irrigations
- Sallyfun Salvia will flower easily for early season requirements

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks	Average Day	Pinch	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing
(12 cm) 5” pot	65° to 68°F (18° to 20°C)		Day Neutral	Soil EC 1.0 - 1.5 pH 5.8 to 6.2
6 to 7 wks				
(15 cm) 6 ½”				
7 to 9 wks				
(20 cm) 8”	3 plants per container			
7 to 9 wks				
(25 cm) 10”	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrips	METHIOCARB,ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide for height control / Paclobutrazol drench for finishing
- Scout for Aphids, Leafminors and Whiteflies

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.8 to 6.2	1.0 to 1.2	68 to 72 F (20 to 22 C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm as needed	Drench fungicide after transplant	
Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68 to 72 f (20 to 22 C)	feed 100 to 150 ppm		Paclobutrazol @ 1 to 2 ppm drench to finish		Paclobutrazol drench when flower bud emerge

For additional technical growing information:

SALLYFUN™ [CLICK HERE](#)



www.danzigeronline.com |