

TOUCH™

Scaevola aemula



TOUCH™
Indigo

- The White Touch™ variety remains pure white even in low temp
- White Touch™ and Blue Touch™ are great for baskets and combos
- Mature height and width is 30 cm x 20 cm (12 x 8 inches)

More varieties in the TOUCH™ lineup:



Best Seller

TOUCH™
Blue



Best Seller

TOUCH™
White



TOUCH™
Baby Pink

For additional technical growing information:

TOUCH™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |



TOUCH™

Scaevola aemula



PROPAGATION TIPS

- Stick on priority – Number 4 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

(10 - 16")
25 - 41 cm



Height

(6 - 10")
20 - 25 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Recommended	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° - 70° (20° - 21° C)		Weeks 3-5 100 to 150 ppm N	Apply broad spectrum Week 2



Misting specification - Remove from mist as soon as possible
Do not over-mist or saturate soil - Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 2000 to 2500 ppm as required
- Pinch is recommended week 4 or at transplant

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist	Not required		Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	72 / 74 F (22 / 23 C)	Feed 100 ppm to 150 ppm	Not required			
Week 4	5.8 to 6.2	1.0 to 1.2	70 F (21 C)	Feed 100 to 150 ppm	Not required			
Week 5	5.8 to 6.2	1.0 to 1.2	70 F (21 C)	Feed 100 to 150 ppm	Not required	Daminozide as needed		Daminozide @ 2000 to 2500 ppm

For additional technical growing information:

TOUCH™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |



TOUCH™

Scaevola aemula



FINISHING TIPS

- Keep the temperatures moderately warm to allow plants to continue to develop (average daily temperature at 68 to 70 F (20 to 21 C))

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 68° - 70° (20° - 21° C)	Pinch –Yes Night interruption or extended day- Optional	150 - 200 ppm N	Spray Daminozide @ 2500 to 3750 ppm
8 to 9 wks (15 cm) 6 ½"			Soil EC 1.0 - 1.2 pH 5.8 to 6.2	Paclobutrazol drench @ 1 to 2 ppm
8 to 9 wks				
(20 cm) 8"	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Paclobutrazol drench application at 1 to 2 ppm for finishing if required
- Drench a broad spectrum fungicide after transplant
- Scout for Aphids and Fungus Gnats and Mites

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.8 to 6.2	1.0 to 1.2	68 to 70 F (20 to 21 C)	feed 100 to 150 ppm		B-9 as needed	Spray fungicide after transplant	Spray Daminozide @ 2500 to 3750 ppm
Week 6 to 13	5.8 to 6.2	1.0 to 1.2	68 to 70 F (20 to 21 C)	feed 100 to 150 ppm		Paclobutrazol Drench if required		Paclobutrazol drench @ 1 to 2 ppm

For additional technical growing information:

TOUCH™ [CLICK HERE](#)

