

# Lollies™

*Argyranthemum*


- Naturally compact mounded habit
- Uniform series of 5 bright colors
- Early blooming
- Well branched



Mounded



30-36 cm  
12-14 in



30-36 cm  
12-14 in



12-15 cm  
Gal, Patio Pots



 DANZIGER

# The LOLLIES™ Series

NEW  
25



**LOLLIES™**  
Buttermint Imp.  
AR-19-5326



**LOLLIES™**  
Pink Pez

Best Seller



**LOLLIES™**  
Marshmallow




**LOLLIES™**  
Berry Gummy



**LOLLIES™**  
White Chocolate



## *Argyranthemum frutescens*



Stick on priority – Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21°-22°C)	NO	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65°-68° (18°-20° C)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 to 6.0	0.80	72°-74° F (22°-23° C)	50 ppm N in mist				Day of sticking
Week 2	5.8 to 6.0	.80 to .90	72°-74° F (22°-23° C)	50 ppm N in mist				Second fungicide app
Week 3	5.8 to 6.0	.90 to 1.0	Cool to 68°- 70° F (20°-21° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.8 to 6.0	1.0 to 1.2	65°-68° F (18°-20° C)	feed 100 to 150 ppm		Daminozide as needed		



**Irrigation specification** - Remove from mist as soon as possible



Argyranthemum respond to long days for flowering. Light in propagation to 12-hour daylength for optimal growth and to hasten crop time. Daminozide at 750 ppm week 3.

## *Argyranthemum frutescens*

- Pinch is optional
- Cool temperature finishing for best quality

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
7 to 8 wks (12 cm) 5" pot 1 ppp	Average Day 68°-72°F (20°-22°C)	Pinch- Optional  Night interruption or extended day- Optional	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½" pot 1 ppp	Weeks 3 – 5 65°-68° (18° - 20° C)		Soil EC 1.0-1.2 pH 5.5 to 6.0	

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.7 to 6.0	1.0 to 1.2	65°-68° F (18°-20° C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
<b>Week 5 to 13</b>	5.7 to 6.0	1.0 to 1.2	65°-68° F (18°-20° C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB



- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- PGR - Spray Daminozide at 750 to 2000 ppm on young finished plants  
Paclobutrazol drench application at 1 ppm for finishing if required
- Maintain good airflow and allow plants to dry before nightfall
- Scout for Thrip on regular basis