

FRESCO™

Brachyscome angustifolia

- Compact plant with a mounded habit
- Early and continuous flowering
- Thick stems that are easy to stick
- Ideal for small pots and hanging baskets




Mounded



12-15
Gal, Patio
Pots



25-30 cm
8-10 in



10-15 cm
4-6 in



Hanging
Basket



Fresco™ Candy

The FRESCO™ Series




Fresco™ Candy



Fresco™ Purple




Brachyscome angustifolia




Stick on priority – Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° (18° C)		Weeks 3-5 75 to 100 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	72° / 74° F (22° / 23° C)	50 ppm N in mist				Day of sticking
Week 2	5.6 to 6.2	.80 to .90	72° / 74° F (22° / 23° C)	50 ppm N in mist		750 ppm Daminozide		Second fungicide app
Week 3	5.6 to 6.2	.90 to 1.0	Cool to 68° / 70° F (20° / 21° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.6 to 6.2	.90 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm		Daminozide as needed		

 – **Irrigation specification** - Remove from mist as soon as possible

-  – High light and warm temperatures will maintain a good proportion of flowers to foliage.
- Maintain lower EC levels for best growth and flower size since higher EC will hinder plant quality.



Brachyscome angustifolia

- Pinch week 3 in the liner or at transplant
- Monitor EC levels so they are not elevated
- Do not over irrigate in the early stages of plant growing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 8 wks (12 cm) 5" pot	Average Day 68° to 72°F (20° to 22°C)	Pinch No lighting required for flowering	100 - 150 ppm N	In liner and finish stages
8 to 10 wks (15 cm) 6 ½"	Weeks 3 – 5 61° - 68° (16° - 20° C)		Soil EC 0.8 - 1.0 pH 5.6 to 6.2	

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.6 to 6.2	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
Week 5 to 13	5.6 to 6.2	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	Not Required			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Avoid covering plant meristem at transplant
- Additional Iron (Fe) or Manganese (Mn) may be required, depending on water quality
- PGR - Spray Daminozide at 750 to 2000 ppm on young & finished plants
- Maintain good airflow and allow plants to dry before nightfall