

# EYECONIC™

## *Calibrachoa*

- Strong contrast in the eye color
- Controlled semi trailing habit with huge flowers
- Tested for pH sensitivity
- Great for combinations
- Blooms in a 10 - hour day



Semi  
trailing



30-41 cm  
12-16 in



30-36 cm  
12-14 in



12-15  
Gal, Patio  
Pots



Hanging  
Basket



EYECONIC™ Purple

 DANZIGER

# The EYECONIC™ Series



EYECONIC™ Cherry Blossom Imp.



EYECONIC™ Orange



EYECONIC™ Com. Sunset



EYECONIC™ Apricot



EYECONIC™ Peach



EYECONIC™ Pink




EYECONIC™ Purple



EYECONIC™ Strawberry



## Calibrachoa hybrida



Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.2 / 5.8	0.80	72° / 74° F (22° / 23° C)	50 ppm N in mist			Day of sticking	
Week 2	5.2 to 5.8	.80 to .90	72° / 74° F (22° / 23° C)	50 ppm N in mist			Second fungicide app	
Week 3	5.2 to 5.8	.90 to 1.0	Cool to 68° / 70° F (20° / 21° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.2 to 5.8	1.0 to 1.2	65° to 68° F (18° to 20° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 5	5.2 to 5.8	1.0 to 1.2	65° to 68° F (18° to 20° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		



– **Irrigation specification** - Remove from mist as soon as possible



- Application of K-IBA can hasten rooting.
- Light in propagation for optimal growth and to hasten crop time- as Calibrachoa can be slower to root and develop a plant structure under short days and more-over under low light conditions.
- Daminozide at 1200 to 1500 ppm day of sticking.



## Calibrachoa hybrida

- Pinch is optional
- Quality improves when finished with cool temperatures

Average Time (from liners)	Temperature	Pinch/Daylength Modification	Fertilization	Plant Growth Regulator
7 to 8 wks (12 cm) 5" pot 1 ppp	Average Day 65° - 68° (18° - 20° C)	No pinch necessary	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½" pot 1 ppp	65° - 68° (18° - 20° C)	No lighting necessary	Soil EC 1.0 - 1.2 pH 5.5 to 6.0	
8 to 9 wks (20 cm) 8" pot 3 ppp				
8 to 9 wks (25 cm) 10" pot 3 ppp				

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 6 Transplant</b>	5.6 to 6.0	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
<b>Week 6 to 12</b>	5.6 to 6.0	1.0 to 1.2	65° to 68° f (18° to 20° C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB



- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- Scout for Aphids and Thrip on regular basis