

COREOPSIS

Verticillate, grandiflora & hybrida

For additional information and availability:



Stick on priority – Number 3 out of 4 categories

PROPAGATION TIPS

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 18° - 20° C (65° - 68° F)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2



– **Irrigation specification** – Remove from mist as soon as root emerges



– Spray adjuvant the day of sticking. Moderate moisture for the rooting media.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	21 / 22 C	50 ppm N in mist	Required 12 hr daylength		Day of sticking	
Week 2	5.8 to 6.5	.80 to .90	21 / 22 C	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.5	.90 to 1.0	Cool to 20 / 21 C	Feed 100 ppm N				
Week 4	5.8 to 6.5	.90 to 1.0	20 to 21 C	Feed 100 ppm N				
Week 5	5.8 to 6.5	.90 to 1.0	18 to 21 C	Feed 100 ppm N		Optional		

COREOPSIS

For additional information and availability:

[COREOPSIS](#)



Verticillate, grandiflora & hybrida

- Pinch required for quality finishing (pinch leaving 4 to 5 nodes)
- Daminozide spray application @ 2000 to 2500 ppm
- Cold beneficial (weeks of night temperatures below 55 F)

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 10 wks	Average Day	Pinch required/ Obligate long day	100 - 150 ppm N	Daminozide spray application @ 2000 to 2500 ppm
15 cm (1 gallon)	(18° to 20° C)			
8 to 10 wks	3 plants per container		Soil EC 0.8 - 1.0	
20 cm (2 gallon)			pH 5.8 to 6.5	

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB

FINISHING TIPS



- Long Day Flowering Response – June through October
- Natural foliar disease resistance
- Maintain good airflow and allow plants to dry between irrigations
- Drench after transplant for Rhizoctonia and Pythium / Phytophthora

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.5	0.8 to 1.0	20° to 21° C	feed 100 to 150 ppm			Drench fungicide after transplant	Control of Pythium & Rhizoctonia
Week 6 to 13	5.8 to 6.5	0.8 to 1.0	18° to 20° C	feed 150 to 200 ppm			Spray fungicide in finishing protocols	Control of Bacterial Leaf Blights