

# GOLF™

## Craspedia


- A true attention grabber
- Unique yellow blooms
- Easier rooting, even in northern climates
- Can be easily cut and put in a vase inside the house
- Well suited for combinations



**Upright**



**20-25 cm**  
**8-10 in**



**36-46 cm**  
**14-18 in**



**Heat**  
**tolerant**



**Pollinator**  
**Friendly**



**Landscape**



**15-18 cm**  
**Gal**



**PLANT OF THE YEAR 2022**  
Pillnitz, North Germany



**DANZIGER**

# GOLF™ BEAUTY IMP.





Stick on priority – Number 1 out of 4 categories  
 Don't store cuttings and plugs in cold store

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 / 6 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Suggested	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 – 5/6 66° - 68° (19° - 20° C)		Weeks 3-5 100 to 150 ppm N	Spray fungicide to control Botrytis and bacteria Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.4	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Optional		Day of sticking	Additional lighting will hasten flowering
Week 2	5.8 to 6.4	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist	Optional		Second fungicide app	Fungicide to control Botrytis!
Week 3	5.8 to 6.4	.90 to 1.0	Cool to 68° / 70° F (20° / 21° C)	Feed 100 ppm to 150 ppm	Optional	Daminozide as needed		
Week 4	5.8 to 6.4	1.0 to 1.2	65° to 68° F (18° / 20° C)	Feed 100 ppm to 150 ppm	Optional	Daminozide as needed		
Week 5	5.8 to 6.4	1.0 to 1.2	65° to 68° F (18° / 20° C)	Feed 100 ppm to 150 ppm	Optional	Daminozide as needed		



– Irrigation specification – Reduce mist shortly after sticking the cuttings (day 2 or 3).



- Dip cut ends of cuttings in rooting hormone prior to sticking or Spray application of K-IBA- This will enable more uniform rooting.
- Daminozide at 1200 to 1500 ppm day of sticking. Pinch plants when they begin to elongate to create branching at base of plant.

*Craspedia spp*

- Pinch week 3 in liner or at transplant
- Plant liners into new soil to support plants at maturity
- Don't store (cuttings and plugs in cold store)

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.8 to 6.2	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm		B-9 as needed	Spray fungicide after transplant	
Week 6 to 14	5.8 to 6.2	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm		Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench if required

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 9 wks (12 cm) 5" pot 1 ppp	Average Day 68° to 72°F (20° to 22°C)	Pinch –Yes  Additional lighting not required	100 - 150 ppm N	In liner and finish stages
8 to 10 wks (15 cm) 6 ½" 1 ppp	65° - 68° (18° - 20° C)		Soil EC 1.0 - 1.2 pH 5.8 to 6.4	

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrip	METHIOCARB,ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB




- PGR - Spray Daminozide at 2000 to 2500 ppm for finishing plants
- Paclobutrazol drench application at 1 to 2 ppm for finishing if required
- Drench broad spectrum fungicide after transplant
- Scout for Aphids on regular basis



Craspedia GOLF™ - Photos from commercial production at Europe, 2022



12cm		
11cm		
10cm		
9cm		
8cm		
7cm		
6cm		
5cm		
4cm		
3cm		
2cm		
1cm		

CRASPEDIA POT

- Golf Beauty

Nota: Cosechar de la base de la planta.