

TRINITY™

Diascia

- Large flowers
- Thicker green leaf with good shelf life
- Exceptional weather tolerance, hot and cold
- Holds flowers longer for improved shelf life
- Grows into a small bush
- Works well in combinations



Upright



32-35 cm
12-14 in



32-38 cm
12-16 in



Heat
Tolerant



12-15
Qt, Gal




Landscape



Pollinator
Friendly



Spring /
Summer
Flowering



TRINITY™ Sunset

 DANZIGER

The TRINITY™ Series

NEW
26



TRINITY™ Salmon
DI-22-3282

NEW
26



TRINITY™ Pink
DI-22-3297



TRINITY™ Grace
DI-20-3106



TRINITY™ Sunset

Diascia barberae

Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° - 70° (20° - 21° C)		Weeks 3-5 100 to 150 ppm N	Apply broad spectrum Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72° / 74° F (22° / 23° C)	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72° / 74° F (22° / 23° C)	50 ppm N in mist	Not required		Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	72° / 74° F (22° / 23° C)	Feed 100 ppm to 150 ppm	Not required			
Week 4	5.8 to 6.2	.90 to 1.0	Cool to 68° / 70° F (20° / 21° C)	Feed 100 to 150 ppm	Not required	Daminozide as needed		Pinch
Week 5	5.8 to 6.2	0.9 to 1.0	68° F (20° C)	Feed 100 to 150 ppm	Not required	Daminozide as needed		



- **Misting specification** - Remove from mist as soon as possible
- Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 1500 to 2000 ppm
- Pinch is recommended week 4 or at transplant



Diascia barberae

- Pinch week 4 in liner or at transplant
- Plant liners above new soil – not to bury plant meristem

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 9 wks	Average Day	Pinch –Yes Day Neutral	100 – 150 ppm N	In liner and finish stages
(12 cm) 5" pot 8 to 10 wks	65° - 68° (18° - 20° C)		Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
(15 cm) 6 ½"	65° - 68° (18° - 20° C)			

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	0.9 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm		Daminozide as needed	Spray fungicide after transplant	
Week 6 to 12	5.8 to 6.2	0.9 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm				Do not apply Paclobutrazol



- PGR - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Paclobutrazol – Do no apply
- Spray broad spectrum fungicide after transplant
- Scout for Aphids and Thrip on regular basis

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB