



# DuraBella®

## Long lasting beauty

Having the right ingredients is the first step in creating a successful recipe, and DURABELLA® is exactly that.

Our teams in the US and Europe come together with a shared passion for excellence, assembling recipe to suit their market's taste and create the perfect blend of varieties. This ensures that everyone can enjoy the beauty of DURABELLA® in their own unique style. These combinations are selected and tested for great performance, resulting in stunning displays that last longer for the consumer.




### Rose Chateau™

Petunia Littletonia™ White Grace

Petunia Capella™ Fuchsia Lace

Verbena Vanessa™ Bicolor Light Pink





Stick on priority - Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks Average for each Genera	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 20° - 21° C (68° - 70° F)		Weeks 3-5 100 ppm N	Apply broad spectrum Week 2



- **Misting specification** - Remove from mist as soon as possible
- Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 2000-2500 on genera requiring control.
- Pinch is recommended week 4 or at transplant on items requiring pinch

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.1	0.80	72 / 74 F	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.6 to 6.1	.80 to .90	72 / 74 F	50 ppm N in mist	Not required		Second fungicide app	
Week 3	5.6 to 6.1	.90 to 1.0	70 / 72 F	Feed 50 ppm to 100 ppm	Not required			
Week 4	5.6 to 6.1	1.0 to 1.2	68 F	Feed 100 to 150 ppm	Not required			
Week 5	5.6 to 6.1	1.0 to 1.2	68 F	Feed 100 to 150 ppm	Not required	Daminozide on items requiring		



- Keep the temperatures moderately warm to allow plants to continue to develop (average daily temperature at 65° F (18° C)
- Plant components close to center of basket to allow them to mix

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (15 cm) 6.5" pot	Average Day 65° ( 18° C )	Pinch – Yes on some items	150 - 200 ppm N	In liner and finish stages on items requiring control
8 to 9 wks (20 cm) 10" or larger	65° ( 18° C )		Soil EC 0.8 - 1.1 pH 5.6 to 6.1	

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB



- PGR – Spot Spray Daminozide at 2500 ppm on components requiring control
- Paclobutrazol drench application at 1 to 2 ppm for finishing if required
- Spray broad spectrum fungicide / drench after transplant when roots are established
- Scout for Aphids, Thrips and Whiteflies on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.6 to 6.1	0.8 to 1.1	65° F	feed 100 to 150 ppm		B-9 as needed	Drench fungicide after transplant	
<b>Week 6 to 13</b>	5.6 to 6.1	0.8 to 1.1	65° F	feed 100 to 150 ppm		Paclobutrazol Drench at 1 to 2 ppm		Drench Paclobutrazol when container is 85% of size for finishing