# CORABELLS<sup>TM</sup> Fuchsia hybrida

- Compact series of Fuchsia
- Early spring flowering ideal for filtered sun locations
- Mature height and width is
- 25 cm x 46 cm (10 x 18 inches)




Early Spring / Spring Flowering



25 - 30 cm (10 - 12")



41 - 46 cm (16 - 18")



## The **CORABELLS™** series



**CORABELLS™** Rose Purple



**CORABELLS™** Red White



**CORABELLS™** Dark Eyes



#### **CORABELLS™**

#### Fuchsia hybrida



Stick on priority – Number 4 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide	
5 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking	
	Weeks 3 - 5 18° - 20° C (65° - 68° F)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2	



- Irrigation specification - Remove from mist as soon as root emerges



 Spray adjuvant the day of sticking. Spray broad spectrum fungicide to prevent Botrytis day of sticking and week 2

Rooting	рН	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	21 / 22 C	50 ppm N in mist			Day of sticking	
Week 2	5.6 to 6.2	.80 to .90	21 / 22 C	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.5	.90 to 1.0	Cool to 20 / 21 C	Feed 100 ppm N				
Week 4	5.8 to 6.5	.90 to 1.0	18 to 20 C	Feed 100 ppm N				
Week 5	5.8 to 6.5	.90 to 1.0	18 to 20 C	Feed 100 to 150 ppm		Should not be required		



### **CORABELLS™**

#### Fuchsia hybrida

- Pinch at transplant
- Well branched and naturally compact plant habit
- Scout for Aphids, Thrips and Whitefly

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator	
7 to 8 wks	Average Day		150 - 200 ppm		
15 cm (1 gallon) 1 ppp	(16° to 18°C)	Pinch required/ possible a second	N N	Should not be required	
8 to 10 wks	3 plants per	pinch if grown in larger containers	Soil EC 0.8 – 1.2		
28 cm (HB) 3 ppp	container	larger containers	pH 5.8 to 6.5		

Finishing	pН	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	0.8 to 1.0	18 to 20 °C	feed 100 to 150 ppm			Drench fungicide after transplant	Control of Pythium & Phytophthora
Week 6 to	5.8 to 6.2	0.8 to 1.2	16° to 18°C	feed 150 to 200 ppm			Spray fungicide in finishing protocols	Control of Bacterial Leaf Blights



- Maintain good airflow and allow plants to dry between irrigations
- High quality plants are grown with moderate/cool temperatures and high light
- Drench after transplant for Rhizoctonia and Pythium / Phytophthora

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS					
resis	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD					
	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN					
Diseases	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM					
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB					



CORABELLS™