

MENOR™

Nemesia hybrida

- Covered in multicolored blooms
- Compact habit, perfect for 6 packs
- Flower early spring



25-30 cm
10-12 in



16-21 cm
6-8 in



10-12 cm



Mounded



MENOR™

 DANZIGER

The MENOR™ Collection



Menor™ Dual Purple




Menor™ Dual Bordeaux



Menor™ Dual Magenta



Menor™ Dual Violet



Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65° - 68° (18° - 20° C)		Weeks 3-4 100 to 125 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6/6.0	0.80	72°/74°F (22°/23°C)	50 ppm N in mist		Daminozide as needed	Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.0	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm to 150 ppm				
Week 4	5.6 to 6.0	1.0 to 1.2	65° to 68°F (18° to 20°C)	Feed 100 ppm to 150 ppm		Daminozide as needed		



– **Irrigation specification** – Reduce mist as soon as possible



Apply broad spectrum fungicide the day of sticking to control leaf disease. For the more vigorous varieties, apply Daminozide spray at 2000 ppm week 3.



Nemesia Hybrida

- Keep the soil evenly moist, avoiding overly wet conditions, especially during cooler growing periods. Prolonged wet soil can lead to root and foliage issues.
- Finish plants at cool temperatures for the best quality and vibrant flower color.

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 65°-68 F (18°-20° C)	Pinch – Recommended	100 - 200 ppm N Soil EC 1.0 - 1.2 pH 5.6 to 6.0	Daminozide spray application at 2500 to 3750 ppm
8 to 9 wks (15 cm) 6 ½"	1 ppp			
8 to 9 wks (20 cm) 8 "	3 ppp			Paclobutrazol drench application at 2 ppm

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.6 to 6.0	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 150 to 200 ppm	Not required	Daminozide as needed 2500 ppm	Drench fungicide after transplant	
Week 6 to 12	5.6 to 6.0	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 150 to 200 ppm	Not required	Paclobutrazol Drench at 2 ppm		Paclobutrazol 2 ppm drench when plants reach 85% of desired size

FINISHING TIPS



- **PGR** – Menor series is naturally compact, therefore PGR application is optional. If needed Daminozide application of up to 2500 ppm on young finished plants is possible.
- **Maintain good airflow** and allow plants to dry before nightfall
- **Scout** for Aphids, Thrips, Mites and Whitefly