



## *Petunia*

- Unique bloom coloration
- Green ruffled flower edges
- Drought tolerance
- Excellent garden performance
- Adds an interesting element to combinations
- Blooms early enough for any market



Compact



Landscape



26-46 cm  
14-18 in



30-36 cm  
12-14 in




12-15  
Gal, Patio Pot



Hanging  
basket



AMAZONAS™ Plum Cockatoo



Stick on priority – Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm	Day of sticking & Week 2



– **Irrigation specification** – Reduce mist as soon as possible



The series is well branched so pinching is not need for a quality finish.  
Daminozide at 1500 ppm the day after sticking. Week 4 – Daminozide @ 2500 ppm / Uniconazole @ 10 ppm tank mix if plants are aggressively growing.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6/6.0	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Additional lighting is optional depending on region	Daminozide day after sticking	Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.0	.90 to 1.0	Cool to 68°F (20°C)	Feed 100 ppm to 150 ppm				
Week 4	5.6 to 6.0	1.0 to 1.2	65° to 68°F (18° to 20°C)	Feed 100 ppm to 150 ppm				

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks	Average Day	No pinch necessary	150 - 200 ppm N	Finish stage Daminozide @ 1500 to 2500 ppm /
(12 cm) 5” pot	68°-72° F (20°-22° C)	Additional lighting is optional depending on region	Soil EC 1.2 - 1.5 pH 5.6 to 6.0	
8 to 9 wks				
(15 cm) 6 ½”				
8 to 10 wks				
(20 cm) 8”	3 plants per container			
8 to 10 wks				
(25 cm) 10”	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.6 to 6.0	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	N/A	Daminozide spray applications	Spray fungicide after transplant	
Week 5 to 15	5.6 to 6.0	1.2 to 1.5	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	N/A	Paclobutrazol Drench at 1 ppm if needed		



- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- PGR – Amazonas genetics are naturally more compact. Spray Daminozide at lower rates if needed.