

# Petunia **splash** DANCE

- Unique speckled pattern
- Tested and trialed to hold its pattern into the hit
- Blooms early enough for any market



**Mounded**



**46-61 cm  
18-24 in**



**25-30 cm  
10-12 in**




**Hanging  
Basket**



**Landscape**



**12-18  
Qt, Gal,  
Patio Pot**



<https://www.youtube.com/watch?v=i1AO2J4M1tQ>



**SPLASH DANCE™ Bolero Blue**



# The **SPLASH DANCE®** series

**Splash Dance® Magenta Mambo**



**Splash Dance®  
Bolero Blue**



**Splash Dance®  
Fuchsia Flamenco**



**Splash Dance®  
Moon Walk**



**Splash Dance®  
Rumba Rose**



**Splash Dance®  
Violet Vogue**




**Splash Dance®  
Purple Polka**



# SPLASH DANCE™

*Petunia Hybrida*

For additional information and availability: [SPLASH DANCE™](#) 



Stick on priority – Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm	Day of sticking & Week 2



**Irrigation specification** – Reduce mist as soon as possible



The series is well branched so pinching is not need for a quality finish.  
 Daminozide at 1500 ppm the day after sticking.  
 Week 4 – Daminozide @ 2500 ppm spray application


Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6/6.0	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Additional lighting is optional depending on region	Daminozide day after sticking	Day of sticking	
Week 2	5.6/6.0	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6/6.0	.90 to 1.0	Cool to 68°F (20°C)	Feed 100 ppm to 150 ppm				
Week 4	5.6/6.0	1.0 to 1.2	65° to 68°F (18° to 20°C)	Feed 100 ppm to 150 ppm		Daminozide @ 2500 spray application		

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks	Average Day	No pinch necessary  Additional lighting is optional depending on region	150 - 200 ppm N	Finish stages Daminozide @ 2500 to 3200 ppm
(12 cm) 5” pot	68°-72° F (20°-22° C)			
8 to 9 wks			Soil EC 1.2 - 1.5 pH 5.6 to 6.0	
(15 cm) 6 ½”				
8 to 10 wks				
(20 cm) 8”	3 plants per container			
8 to 10 wks				
(25 cm) 10”	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.6 to 6.0	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	N/A	Daminozide/ Uniconazole tank mix if required	Spray fungicide after transplant	
Week 6 to 13	5.6 to 6.0	1.2 to 1.5	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	N/A	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

- 
- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
  - PGR - Spray Daminozide at 2,500 to 3,200 ppm and / or Paclobutrazol drench application at 1 to 2 ppm for finishing if required. The most vigorous varieties are Bolero Blue, Purple Polka and Violet Vogue