

# MEGA PAZZAZ™

Portulaca oleracea


- Big flowers (MEGA blooms) on a moderate semi trailing habit
- Bright colors grab attention at retail and in the garden
- Strong option for heat/drought tolerant programs
- Well suited for combinations



-Semi  
Trailing



25-30 cm  
10-12 in



41-51 cm  
in16-20



Heat  
tolerant



Pollinator  
Friendly



Landscape



Ground  
cover



12-15  
Qt, Gal



Hanging  
basket



MEGA PAZZAZ™ Mango Twist



# The MEGA PAZZAZ™ Series

NEW  
25



**MEGA PAZZAZ™ Papaya Twist**  
PT-21-1911

NEW  
25



**MEGA PAZZAZ™ Tropical Twist**  
PT-21-1833



**MEGA PAZZAZ™ Dark Pink**



**MEGA PAZZAZ™ Gold**



**MEGA PAZZAZ™ Pink Twist**

NEW  
25



**MEGA PAZZAZ™ Fuchsia**  
PT-21-1836



**MEGA PAZZAZ™ Mango Twist**



**MEGA PAZZAZ™ Orange**



**MEGA PAZZAZ™ Purple**




**MEGA PAZZAZ™ Red**

# PAZZAZ NANO™

*Portulaca oleracea*

For additional information and availability:

MEGA PAZZAZ™



Stick on priority – Number 1 out of 4 categories

PROPAGATION TIPS

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8/6.2	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8/6.2	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8/6.2	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm to 150 ppm				
Week 4	5.8/6.2	1.0 to 1.2	68° to 70°F (20°/21°C)	Feed 100 to 150 ppm		Daminozide as required		



– **Irrigation specification** – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking. Depending on growth, pinch week 3 or at transplant to allow improved branching.

## *Portulaca oleracea*

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks (12 cm) 5" pot	Average Day 68°-72° F (20°-22° C)	Pinch 1 to 2 times  Additional lighting will hasten flowering under short days	150 - 200 ppm N	Daminozide @ 2500 ppm or tank mix of Daminozide / Uniconazole for finishing
6 to 7 wks (15 cm) 6 ½"			Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
7 to 9 wks				
(20 cm) 8"	3 plants per container			

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB / Triflumizole

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 4 Transplant</b>	5.8 to 6.2	1.0 to 1.2	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	
<b>Week 5 to 12</b>	5.8 to 6.2	1.2 to 1.5	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm or tank mix of Daminozide / Uniconazole for finishing		



- **PGR** - Spray Daminozide / Uniconazole tank mix to control height
- Scout for Aphids, Thrips and Whiteflies