

# PAZZAZ NANO™

#### Portulaca oleracea

- Naturally controlled habit
- Early flowering
- Big flowers that stay open longer
- Strong option for heat/drought tolerant programs
- Performs well in combinations
- Excellent option for pollinator programs





# The **PAZZAZ NANO™** series



PAZZAZ<sup>™</sup> NANO Scarlet Twist PT-21-1933



PAZZAZ<sup>™</sup> NANO Deep Orange



PAZZAZ<sup>™</sup> NANO Candy Pink



PAZZAZ<sup>™</sup> NANO Hot Pink



PAZZAZ™ NANO Gold



PAZZAZ<sup>™</sup> NANO Fuchsia Imp.



Mango



PAZZAZ<sup>™</sup> NANO Orange



PAZZAZ™ NANO Orange Twist



PAZZAZ<sup>™</sup> NANO Purple PT-21-1852



PAZZAZ<sup>™</sup> NANO Tropical Punch



PAZZAZ™ NANO Yellow



PAZZAZ<sup>™</sup> NANO Yellow Twist



## PAZZAZ NANO™

Portulaca oleracea





#### Stick on priority – Number 1 out of 4 categories

	verage Fime	Temperature	Hormone	Fertilization	Fungicide			
4 \	weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking			
		Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2			



 Irrigation specification – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking.

Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	рН	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8/6.2	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8/6.2	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8/6.2	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm to 150 ppm				
Week 4	5.8/6.2	1.0 to 1.2	68° to 70°F (20°/21°C)	Feed 100 to 150 ppm		Daminozide as required		



### PAZZAZ NANO™

#### PAZZAZ NANO™

### Portulaca oleracea

FINISHING TIPS

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator			Aphids		FTAMIPRID					
5 to 6 wks	Average Day	Pinch Additional lighting will hasten flowering under short days	150 - 200	Daminozide @ 2500 ppm for finishing if required Diseases	Pests		Thrips		ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD					
(12 cm) 5" pot	68°-72° F (20°-22° C)		ppm N		Diseases		Botrytis Pythium		CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN PROPAMOCARB / MEFENOXAM					
6 to 7 wks			Soil EC 1.0 - 1.2 pH 5.8 to 6.2		Pł	Phytophthora Photophthora   Rhizoctonia AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB / T					Triflumizole			
(15 cm) 6 ½"					Finishing	рН	EC	Temp	Feed	Light	PGR	Fungicide	Comment	
7 to 9 wks										8				
(20 cm) 8″	3 plants per container				Week 4 Transplant	5.8 to 6.2	1.0 to 1.2	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant		
0. :	Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm for finishing							

