

PAZZAZ™

Portulaca oleracea

- Moderate a semi trailing habit that works well for gallon and hanging basket production
- Early flowering
- Strong option for heat/drought tolerant programs
- Performs well in combinations
- Excellent option for pollinator programs



Semi-
Trailing



10 to 12in
25 to 30cm



14 to 16in
36 to 41cm



Heat
Tolerant



12-15
Qt, Pack




Landscape



Pollinator
friendly



Late Spring /
Summer
Flowering



Balanced Beauty- PAZZAZ™

PAZZAZ™ strikes the perfect balance with its naturally medium habit, making it versatile for various garden settings, from hanging baskets to mid-sized containers, offering a splendid range of colors to brighten any spot.



PAZZAZ™ Tangerine



PAZZAZ™ Campino
Twist Pink



PAZZAZ™ Campino
Twist Red



PAZZAZ™
Red Flare Imp.



PAZZAZ™
Yellow Improved



PAZZAZ™
White



Stick on priority – Number 1 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° - 70° (20° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



– **Irrigation specification** – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking. Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8/6.2	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	68° to 70°F (20°/21°C)	Feed 100 to 150 ppm		Daminozide as required		

Portulaca oleracea

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks (12 cm) 5" pot	Average Day 68°-72° F (20°-22° C)	Pinch	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing
6 to 7 wks (15 cm) 6 ½"		Additional lighting will hasten flowering under short days	Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
7 to 9 wks				
(20 cm) 8"	3 plants per container			



- PGR - Spray Daminozide control height
- Scout for Aphids, Thrips and Whiteflies

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS						
	Thrips	METHIOCARB,ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD						
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN						
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM						
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB / Triflumizole						
Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 4 Transplant	5.8 to 6.2	1.0 to 1.2	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide at transplant	
Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm for finishing		