

# SALLYFUN™

Salvia frinacea

- Easy to grow
- Good for combos
- Continues flowering into summer
- Excellent choice for pollinator programs
- Works well as an upright element in a combination



Upright



10-18 in  
25-46 cm



12-16 in  
30-41 cm



12-15



Landscape



# The SALLYFUN™ series

## Medium



SALLYFUN™ Bicolor Blue



SALLYFUN™ Blue Ice



SALLYFUN™ Blue Lagoon



STREAM™ Deep Ocean



SALLYFUN™ Pure White

## Large




SALLYFUN™ Blue



SALLYFUN™ Blue Emotion



SALLYFUN™ Snowwhite



Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 18° - 21° C (65° - 70° F)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



– **Irrigation specification** – Reduce mist as soon as possible



- Spray adjuvant at sticking . Daminozide at 2000 ppm the day after sticking.
- Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F	50 ppm N in mist	Netural	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68 / 70 F	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	65 to 68 F	Feed 100 to 150 ppm		Daminozide as required		





- Allow plants to dry between irrigations
- Sallyfun Salvia will flower easily for early season requirements

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks (12 cm) 5" pot	Average Day 65° to 68°F (18° to 20°C)	Pinch Day Natural	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing
6 to 7 wks (15 cm) 6 ½"			Soil EC 1.0 - 1.5 pH 5.8 to 6.2	Paclobutrazol 1 to 2 ppm drench to hold plants
7 to 10 wks (20 cm) 8"	3 plants per container			
7 to 9 wks (25 cm) 10"	3 plants per container			



- PGR - Spray Daminozide for height control / Paclobutrazol drench for holding
- Scout for Aphids, Leafminors and Whiteflies

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 4 Transplant</b>	5.8 to 6.2	1.0 to 1.2	68 to 72 F	feed 100 to 150 ppm		Daminozide @ 2500 ppm as needed	Drench fungicide after transplant	
<b>Week 5 to 12</b>	5.8 to 6.2	1.2 to 1.5	68 to 72 F	feed 100 to 150 ppm		Paclobutrazol @ 1 to 2 ppm drench to finish		Paclobutrazol drench when flower bud emerge