

# BLESSING™

*Scaevola aemula*

- Vigorous semi-trailing habit
- large flowers, open simultaneously for consistent color
- Semi-upright structure ideal for quart and larger pots
- Well-suited for heat-tolerant growing programs.



Semi-  
Upright



10-20 cm  
4-8 in



36-46 cm  
14-18 in



Heat  
Tolerant



12-15  
Patio Pot



Landscape



Hanging  
Basket



Blessing™ series

 **DANZIGER**

# The **BLESSING™** series

NEW  
26



**BLESSING™ Bicolor Rose**  
SC-22-1895



**BLESSING™ Bicolor Lavender**



**BLESSING™ Lavender**



**BLESSING™ Pink**



Stick on priority – Number 4 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Recomm ended	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° - 70° (20° - 21° C)		Weeks 3-5 100 to 150 ppm N	Apply broad spectrum Week 2



- **Misting specification** - Remove from mist as soon as possible
- Do not over-mist or saturate soil - Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 2000 to 2500 ppm as required
- Pinch is recommended week 4 or at transplant

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8/6.2	0.80	72°/74°F (22°/23°C)	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist	Not required		Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	72°/74°F (22°/23°C)	Feed 100 ppm to 150 ppm	Not required			
Week 4	5.8 to 6.2	1.0 to 1.2	70°F (21°C)	Feed 100 ppm to 150 ppm	Not required			
Week 5	5.8 to 6.2	1.0 to 1.2	70°F (21°C)	Feed 100 ppm to 150 ppm	Not required	Daminozide as needed		Daminozide @ 2000 to 2500 ppm

Scaevola aemula

- Keep the temperatures moderately warm to allow plants to continue to develop (average daily temperature at 68 to 70 F (20 to 21 C)

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks	Average Day	Pinch – Yes  Night interruption or extended day-Optional	150 - 200 ppm N	Spray Daminozide @ 2500 to 3750 ppm
(12 cm) 5” pot	68°-70° F (20°-21° C)		Soil EC 1.0 - 1.2 pH 5.8 to 6.2	Paclobutrazol drench @ 1 to 2 ppm
8 to 9 wks				
(15 cm) 6 ½”				
8 to 9 wks				
(20 cm) 8”	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.8 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21° C)	feed 100 to 150 ppm		B-9 as needed	Spray fungicide after transplant	Spray Daminozide @ 2500 to 3750 ppm
Week 6 to 13	5.8 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21° C)	feed 100 to 150 ppm		Paclobutrazol Drench if required		Paclobutrazol drench @ 1 to 2 ppm



- PGR - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Paclobutrazol drench application at 1 to 2 ppm for finishing if required
- Drench a broad spectrum fungicide after transplant
- Scout for Aphids and Fungus Gnats and Mites