

TOUCH™

Scaevola aemula

- Naturally controlled habit
- Abundance of flowers
- Season extender
- Great summer item
- Well suited for combinations



Mounded



20-25 cm
8-10 in



25-41 cm
10-16 in



Heat
tolerant



12-15
Qt, Gal



Landscape



Hanging
basket

Touch™ white

 **DANZIGER**



The TOUCH™ series



TOUCH™ Baby Pink




TOUCH™ Blue



TOUCH™ Indigo



BLESSING™ White

*Scaevola aemula*

Stick on priority – Number 4 out of 4 categories

| Average Time | Temperature | Hormone | Fertilization | Fungicide |
|--------------|--|-------------|------------------------|--|
| 5 weeks | Weeks 1 - 2 70°-72° F (21° - 22°C) | Recommended | Weeks 1-2 50 ppm N | Spray fungicide to control Botrytis and bacteria day of sticking |
| | Weeks 3 - 5 68° - 70° (20° - 21° C) | | Weeks 3-5 150 ppm N | Apply broad spectrum Week 2 |



- **Misting specification** - Remove from mist as soon as possible
- Do not over-mist or saturate soil - Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 2000 to 2500 ppm as required
- Pinch is recommended week 4 or at transplant

| Rooting | pH | EC | Temp | Feed | Light | PGR | Fungicide | Comments |
|---------|---------------|------------|------------------------|-------------------------------|-----------------|-------------------------|----------------------------|-------------------------------------|
| Week 1 | 5.8/6.2 | 0.80 | 72°/74°F (22°/23°C) | 50 ppm N in mist | Not required | | Day of sticking | |
| Week 2 | 5.8 to 6.2 | .80 to .90 | 72°/74°F (22°/23°C) | 50 ppm N in mist | Not required | | Second fungicide app | |
| Week 3 | 5.8 to 6.2 | .90 to 1.0 | 72°/74°F (22°/23°C) | Feed 100 ppm to 150 ppm | Not required | | | |
| Week 4 | 5.8 to 6.2 | 1.0 to 1.2 | 70°F (21°C) | Feed 100 ppm to 150 ppm | Not required | | | |
| Week 5 | 5.8 to 6.2 | 1.0 to 1.2 | 70°F (21°C) | Feed 100 ppm to 150 ppm | Not required | Daminozide as needed | | Daminozide @ 2000 to 2500 ppm |

*Scaevola aemula*

- Keep the temperatures moderately warm to allow plants to continue to develop (average daily temperature at 68 to 70 F (20 to 21 C))

| Average Time (from liners) | Temperature | Pinch/ Daylength Modification | Fertilization | Plant Growth Regulator |
|------------------------------|---|---|------------------------------------|-------------------------------------|
| 6 to 7 wks (12 cm) 5" pot | Average Day 68°-70° F (20°-21° C) | Pinch –Yes | 150 - 200 ppm N | Spray Daminozide @ 2500 to 3750 ppm |
| 8 to 9 wks (15 cm) 6 ½" | | Night interruption or extended day-Optional | Soil EC 1.0 - 1.2 pH 5.8 to 6.2 | Paclobutrazol drench @ 1 to 2 ppm |
| 8 to 9 wks | | | | |
| (20 cm) 8 " | 3 plants per container | | | |

| Pests | Aphids | ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS |
|----------|--------------|--|
| | Thrip | METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD |
| Diseases | Botrytis | CYPRODINIL+FLUDIOXONIL, IPRRODIONE, POLYOXIN |
| | Pythium | PROPAMOCARB |
| | Phytophthora | PROPAMOCARB |

| Finishing | pH | EC | Temp | Feed | Light | PGR | Fungicide | Comments |
|--------------------------|------------|------------|--------------------------------|---------------------|-------|----------------------------------|----------------------------------|-------------------------------------|
| Week 5 Transplant | 5.8 to 6.2 | 1.0 to 1.2 | 68° to 70° F (20° to 21° C) | feed 100 to 150 ppm | | B-9 as needed | Spray fungicide after transplant | Spray Daminozide @ 2500 to 3750 ppm |
| Week 6 to 13 | 5.8 to 6.2 | 1.0 to 1.2 | 68° to 70° F (20° to 21° C) | feed 100 to 150 ppm | | Paclobutrazol Drench if required | | Paclobutrazol drench @ 1 to 2 ppm |



- PGR - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Paclobutrazol drench application at 1 to 2 ppm for finishing if required
- Drench a broad spectrum fungicide after transplant
- Scout for Aphids and Fungus Gnats and Mites