


STREAM COMPACT™

Lobularia hybrida

- Controlled growth in the greenhouse then continues to gain size to fill larger containers
- Floriferous, ball-shaped plant structure
- Highly fragrant, pollinators love it!
- Flowers from early spring through autumn
- Less PGR needed
- Ideal as a filler in mixed containers, hanging basket or window boxes
- Heat and cold tolerant, very versatile



Mounded



cm 10-15
4-6 in



cm 25-46
10-18 in



Heat
Tolerant



12-15
Qt, Gal



Landscape



Ground
Cover



Pollinator
Friendly



Spring /
Summer
Flowering



STREAM™ Compact White

The STREAM COMPACT™ series



STREAM™ Compact Purple



STREAM™ Compact Rose



STREAM™ Compact Violet

Best Seller



STREAM™ Compact White


STREAM™ Compact

For additional information and availability:

STREAM™ Compact



Lobularia hybrida



Stick on priority – Number 1 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2



– **Irrigation specification** – Reduce from mist as soon as possible



– Application of K-IBA can hasten rooting. Daminozide at 1500 to 2500 ppm day of sticking. Lighting in propagation is not necessary. The Lobularia Stream™ Series is well branched and a pinch would only be necessary if grown under lower light levels.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.2 to 5.8	0.80	70°-72° F (21° - 22°C)	50 ppm N in mist			Day of sticking	Feeding through Mist
Week 2	5.2 to 5.8	.80 to .90	70°-72° F (21° - 22°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.2 to 5.8	.90 to 1.0	Cool to 68° - 70° F (20° - 21° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.2 to 5.8	1.0 to 1.2	65° to 68° F (18° to 20° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 5	5.2 to 5.8	1.0 to 1.2	65° to 68° F (18° to 20° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		



Lobularia hybrida

- Pinch is optional
- Cool temperature finishing for best quality

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 65° - 68° (18° - 20° C)	Pinch / Optional	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½"	1 plant per pot		Soil EC 0.8 - 1.0 pH 5.6 to 6.0	
8 to 9 wks				
(20 cm) 8"	3 plants per container			
8 to 9 wks				
(25 cm) 10"	3 plants per container			

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.0	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
Week 6 to 12	5.6 to 6.0	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrips	METHIOCARB,ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Consistent soil moisture levels are best – allow soil to dry slightly between irrigations
- Do not apply chemicals with high tendency to create leaf damage
- Scout for Aphids on regular basis