

MEGA PAZZAZ™

Portulaca oleracea

- Big flowers (MEGA blooms) on a moderate semi trailing habit
- Bright colors grab attention at retail and in the garden
- Strong option for heat/drought tolerant programs
- Well suited for combinations

				
-Semi Trailing	25-30 cm 10-12 in	41-51 cm in16-20	Heat tolerant	Pollinator Friendly
				
Landscape	Ground cover	12-15 Qt, Gal	Hanging basket	



MEGA PAZZAZ™ Mango Twist

The MEGA PAZZAZ™ Series



MEGA PAZZAZ™ Papaya Twist



MEGA PAZZAZ™ Tropical Twist



MEGA PAZZAZ™ Dark Pink

Best Seller



MEGA PAZZAZ™ Gold



MEGA PAZZAZ™ Pink Twist




MEGA PAZZAZ™ Fuchsia



MEGA PAZZAZ™ Mango Twist

Best Seller



MEGA PAZZAZ™ Orange

Best Seller




MEGA PAZZAZ™ Purple



MEGA PAZZAZ™ Red

Portulaca oleracea



Stick on priority – Number 1 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 72°-74° F (22° - 23°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 to 6.2	0.80	72°-74° F (22° - 23°C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72°-74° F (22° - 23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68° - 70° (20° - 21° C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	68° - 70° (20° - 21° C)	Feed 100 to 150 ppm		Daminozide as required		



– **Irrigation specification** – Reduce mist as soon as possible/excess misting will create uneven rooting and delay rooting



This genera is top priority for sticking. Remove plants from box upon receiving allowing air-movement around bags. Daminozide at 1500 ppm the day after sticking. Depending on growth, pinch week 3 or at transplant to allow improved branching.

*Portulaca oleracea*

- Allow plants to dry completely between irrigations
- Maintain warm temperatures allowing plants to develop

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks	Average Day	Pinch 1 to 2 times	100 to 150 ppm N	Daminozide @ 2500 ppm or tank mix of Daminozide / Uniconazole for finishing
(12 cm) 5" pot	68°-72° F (20°-22° C)			
6 to 7 wks	1 plant per pot	Additional lighting will hasten flowering under short days	Soil EC 1.0 - 1.5 pH 5.8 to 6.2	
(15 cm) 6 ½"				
7 to 9 wks				
(20 cm) 8"	3 plants per container			



- PGR - Spray Daminozide / Uniconazole tank mix to control height
- Scout for Aphids, Thrips and Whiteflies

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB / Triflumizole

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Transplant to Week 4	5.8 to 6.2	1.0 to 1.2	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide as needed	Drench fungicide after transplant	
Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68° to 72° F (20° to 22° C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm or tank mix of Daminozide / Uniconazole for finishing		