

SALLYFUN™

Salvia farinacea

- Easy to grow
- Continuous flowering throughout the summer
- Excellent choice for pollinator programs
- Works well as an upright element in combinations



Upright



10-18 in
25-46 cm



12-16 in
30-41 cm



Heat
Tolerant



12-15
Gal,
Patio Pot



Landscape



Pollinator
friendly



The SALLYFUN™ series

Medium



SALLYFUN™ Bicolor Blue



SALLYFUN™ Blue Ice

Best Seller



SALLYFUN™ Blue Lagoon

Best Seller



STREAM™ Deep Ocean



SALLYFUN™ Pure White

Large



SALLYFUN™ Sky Blue



SALLYFUN™ Blue




SALLYFUN™ Blue Emotion



SALLYFUN™ Snowwhite

Salvia farinacea



Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 18° - 20° C (65° - 68° F)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2



- **Irrigation specification** – Reduce mist as soon as possible



- Spray adjuvant at sticking . Daminozide at 2000 ppm the day after sticking.
- Depending on growth, pinch week 3 or at transplant to allow improved branching.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 to 6.2	0.80	21° - 22°C (70°-72° F)	50 ppm N in mist	Netural		Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	21° - 22°C (70°-72° F)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 20° - 21°C (68°- 70° F)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	18° - 20° C (65° - 68° F)	Feed 100 to 150 ppm		Daminozide as required		

Salvia farinacea

- Allow plants to dry between irrigations
- Sallyfun Salvia will flower easily for early season requirements

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
5 to 6 wks	Average Day	Pinch Day Natural	150 - 200 ppm N	Daminozide @ 2500 ppm for finishing
(12 cm) 5" pot	68° to 72°F (20° to 22°C)			Paclobutrazol 1 to 2 ppm drench to hold plants
6 to 7 wks	1 plant per pot		Soil EC 1.0 - 1.5 pH 5.8 to 6.2	
(15 cm) 6 ½"				
7 to 10 wks				
(20 cm) 8"	3 plants per container			
7 to 9 wks				
(25 cm) 10"	3 plants per container			

-
- PGR - Spray Daminozide for height control / Paclobutrazol drench for holding
 - Scout for Aphids, Leafminors and Whiteflies

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTROBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Transplant to Week 4	5.8 to 6.2	1.0 to 1.2	68° to 72°F (20° to 22°C)	feed 100 to 150 ppm		Daminozide @ 2500 ppm as needed	Drench fungicide after transplant	
Week 5 to 12	5.8 to 6.2	1.2 to 1.5	68° to 72°F (20° to 22°C)	feed 100 to 150 ppm		Paclobutrazol @ 1 to 2 ppm drench to finish		Paclobutrazol drench when flower bud emerge